

Research paper

A latent class analysis using the integrated motivational-volitional model of suicidal behaviour: Understanding suicide risk over 36 months



Nikki L. van Eijk ^{a,e,*}, Karen Wetherall ^b, Eamonn Ferguson ^c, Daryl B. O'Connor ^d,
Rory C. O'Connor ^{b,1}

^a 113 Suicide Prevention, Amsterdam, the Netherlands

^b Suicidal Behaviour Research Laboratory, Institute of Health and Wellbeing, University of Glasgow, UK

^c School of Psychology, University of Nottingham, UK

^d School of Psychology, University of Leeds, UK

^e Department of Psychology, Open University, the Netherlands

ARTICLE INFO

Keywords:

Suicide

Integrated motivational-volitional (IMV) model

Latent class analysis

Theory

ABSTRACT

Background: The use of latent class analysis (LCA) to understand suicide risk is often not guided by theoretical frameworks. This study used the Integrated Motivational-Volitional (IMV) Model of Suicidal Behaviour to inform the classification of subtypes of young adults with a suicidal history.

Methods: Data from young adults in Scotland ($n = 3508$) were used in this study including a subgroup of participants ($n = 845$) with a history of suicidality. LCA using risk factors from the IMV model was conducted on this subgroup, and the subgroups and non-suicidal control group were compared. Trajectories of suicidal behaviour over 36 months was compared between the classes.

Results: Three classes were identified. Class 1 (62 %) had low scores on all risk factors, Class 2 (23 %) had moderate scores, and Class 3 (14 %) had high scores on all risk factors. Those in Class 1 had a stable low risk of suicidal behaviour, while those in Class 2 and 3 showed marked variation over time, although Class 3 had the highest risk across all timepoints.

Limitations: The rate of suicidal behaviour in the sample was low, and differential dropout may have impacted the findings.

Conclusions: These findings suggest that young adults can be classified into different profiles based on suicide risk variables derived from the IMV model, which still distinguishes them 36 months later. Such profiling may help determining who is most at risk for suicidal behaviour over time.

1. Introduction

Suicide is a complex phenomenon, accounting for 703,000 deaths each year, including among young people (World Health Organization (WHO), 2021). Studies trying to predict suicide attempts have failed to do so accurately (Franklin et al., 2017). As many of the risk factors that are associated with suicide are highly correlated, and there may be complex interactions between the variables that determine suicide risk, traditional statistical approaches may not be suitable for accurately predicting suicides. Some alternatives have been proposed, such as Latent Class Analysis (LCA). LCA models use a person-centred method to

divide a large group of individuals into smaller homogenous subgroups. Because of the complex interplay of factors that contribute to suicide risk, identifying classes of individuals with certain shared characteristics could lead to a way of classifying risk and improving detection and prevention. Several studies have sought to examine suicide using LCA models and examining a variety of risk and protective factors (Au et al., 2021; Bernanek et al., 2017; Hoogstoel et al., 2021; Logan et al., 2011; Love and Durtschi, 2021; Selby and Gardner, 2015; Thullen et al., 2016; Xiao et al., 2019). These studies have often taken different approaches. Some studies utilise representative population samples, which tend to identify one large control group of healthy individuals, and two or more

* Corresponding author at: Paasheuvelweg 25, 1105 BP Amsterdam, the Netherlands.

E-mail addresses: n.vaneijk@113.nl (N.L. van Eijk), karen.wetherall@glasgow.ac.uk (K. Wetherall), eamonn.ferguson@nottingham.ac.uk (E. Ferguson), d.b.oconnor@leeds.ac.uk (D.B. O'Connor), rory.oconnor@glasgow.ac.uk (R.C. O'Connor).

¹ Author details: Rory C O'Connor, University of Glasgow, Scotland.

groups at varying risk of suicidal ideation or attempts (Jiang et al., 2010; Jung et al., 2019; O'Neill et al., 2018; Rice et al., 2018; Zhu et al., 2019). Whereas other studies have focused specifically on those who have made a suicide attempt (Ginley and Bagge, 2017), or on specific risk groups, such as military veterans (Allan et al., 2020) or those in sexual minority groups (Giano et al., 2020).

A limitation of most of the previous LCA and suicide risk studies is that the choice in the risk factors used to identify the classes have often not been explicitly substantiated. Although some studies highlight theoretical frameworks of suicide risk, such as the Interpersonal Psychological Theory of Suicide (IPTS; Joiner, 2005) or the Integrated Motivational-Volitional model of suicidal behaviour (IMV; O'Connor and Kirtley, 2018), only one study has thus far explicitly used the variables of a model, the IPTS, to identify latent classes in a sample of individuals who had made a suicide attempt (Wong et al., 2020). Wong et al. (2020), using data from a relatively small sample ($N = 227$), found three latent classes that differed from each other on some of the key factors in the IPTS, mainly thwarted belongingness and perceived burdensomeness.

Additionally, most previous studies have employed cross-sectional designs when investigating the relationship between latent classes and past or current suicidal thoughts and feelings. Although, some studies have used a longitudinal design, the follow-up time periods were short (Allan et al., 2020) or the outcome was limited to suicidal thoughts rather than suicide attempts (Wong et al., 2020; Zhu et al., 2019). To date, no studies have investigated the stability of the class assignments over time, which is crucial for the clinical utility of categorising people into different subgroups.

For the current study, we used LCA models derived from a theoretical framework, the IMV model, to identify classes of individuals with a history of suicidal ideation and behaviour and used these classes to predict suicidal behaviour over 36 months. The IMV model builds upon several theories of suicide, such as the IPTS, and provides a framework for understanding how suicide risk develops from the pre-motivational phase (i.e., factors that create vulnerability to suicide), to the motivational phase (i.e., factors that lead to the emergence of suicidal thinking), to the volitional phase (i.e., factors that increase the likelihood of acting upon suicidal thoughts). Evidence has been found for the different phases of the IMV model (Branley-Bell et al., 2019; Lucht et al., 2020; O'Connor et al., 2021; Ordóñez-Carrasco et al., 2020; for an overview see Wetherall et al., 2020), and it has successfully distinguished between those who think about suicide and those who attempt suicide (Dhingra et al., 2015; Wetherall et al., 2018). However, we think that within those groups of people with suicidality, more fine-grained differences exist that we would not be able to find by only categorising them into those who think about suicide and those who attempt suicide. Therefore, using the IMV variables to identify latent subgroups or patterns within the data using LCA could increase our understanding of the characteristics of suicidal people. This method allows us to explore which combinations of risk factors co-occur and how these factors relate to the risk of future suicide attempts.

In this study, we identified latent classes using key variables derived from the IMV model among individuals who have had a history of suicidal thoughts or behaviour. These classes were compared with each other, as well as with a comparison group of individuals with no history of suicidal thoughts or behaviour. Specifically, this study had three aims. The first aim was to identify the best fitting number of latent classes and to identify how they differed on key suicide risk variables. The second aim was to use the latent classes to predict suicide attempts longitudinally. Lastly, the third aim was to identify whether the classes still differed at follow up and which of the IMV and suicide risk variables distinguished between the classes 36 months after baseline. If so, the latter would act as a validation of the predictive value of class structures over time as well as identifying key risk variables.

2. Methods

2.1. Participants and procedure

The data were from the Scottish Wellbeing Study (O'Connor et al., 2018), a nationally representative sample of young adults aged 18–34 years ($n = 3508$) in Scotland. Quota sampling was used based on age, sex and working status (for more details, see O'Connor et al., 2018). After providing written consent, participants completed an hour-long interview in their own homes. Participants completed a range of psychological and social measures that incorporated key aspects of the IMV model. Participants were compensated £25 for their time. Participants who agreed to be contacted again were contacted after 12 months, 24 months, and 36 months, to complete a follow-up questionnaire, which was conducted by email, post, or phone. Participants were compensated with shopping vouchers and entered into a prize draw. Ethical approval was obtained from the Psychology Department's ethics committee at the University of Stirling as well as the US Department of Defense Human Research Protections Office.

2.2. Materials

2.2.1. Factors used in the LCA

A number of risk and protective factors for suicidal behaviour was assessed within the Scottish Wellbeing Study (O'Connor et al., 2018). For the creation of the latent classes, six indicator variables were used. These were depressive symptoms as measured by the Beck Depression Inventory (BDI; Beck et al., 1996), defeat and entrapment scales (Gilbert and Allan, 1998), and current suicidal ideation measured with the Scale for Suicide Ideation (SSI; Beck et al., 1988).

2.2.2. Factors used to test the classes

Other risk and protective factors from the Scottish Wellbeing Study were selected to test the classes. These factors were: social support (ENRICH Social Support Instrument (ESSI)) (Mitchell et al., 2003), impulsivity (Barratt Impulsiveness Scale (BIS-11); Patton et al., 1995), thwarted belongingness and perceived burdensomeness (Interpersonal Needs Questionnaire, INQ; van Orden et al., 2012), acquired capability (Acquired Capability for Suicide Scale (ACSS); van Orden et al., 2008), social perfectionism (from the Multidimensional Perfectionism Scale) (Hewitt and Flett, 1991), problematic alcohol use (CAGE questionnaire, Ewing, 1984), and mental wellbeing (Warwick-Edinburgh Mental Well-being Scale (WEMWBS); Tennant et al., 2007). More details on the measures used can be found in (Wetherall et al., 2018), however, note that each of the measures demonstrated good internal reliability in the current study.

2.2.3. Suicidal ideation and suicide attempts

People's history of suicidal thoughts, attempts and self-harm was assessed with three questions drawn from the Adult Psychiatric Morbidity Survey (McManus et al., 2009): "Have you ever seriously thought of taking your life, but not actually attempted to do so?", "Have you ever made an attempt to take your life, by taking an overdose of tablets or in some other way?", and "Have you ever deliberately harmed yourself in any way but not with the intention of killing yourself?". Responses to these items were "no", "yes" or "would rather not say". Suicidal ideation and suicide attempts were assessed at each follow-up (12, 24 and 36 months), with the first two items used to assess the history of suicidal ideation and suicide attempts, but they were adapted to reflect the last 12 months.

2.3. Data analysis

Preliminary analyses were conducted in SPSS 23 (IBM Corp, 2015) to select only those participants who reported lifetime suicidal ideation and/or a lifetime suicide attempt at baseline for inclusion in the latent

class analysis, with a sample of $n = 867$ participants. Participants with missing values on any of the indicator variables used to create the profiles were also excluded, and the final sample size for the LCA was $n = 845$. A Latent Class Analysis was carried out in R, using the tidyLPA package (Rosenberg et al., 2018). The subgroups for this study were defined by six indicator variables: depressive symptoms, current suicidal ideation, defeat, internal entrapment and external entrapment, as well as impulsivity. Solutions were fit for one to six classes. The best-fitting class solution was decided based on the Akaike's Information Criteria (AIC), Bayesian Information Criteria (BIC), the Bootstrapped Likelihood Ratio test (BLRT) and entropy.

The full sample ($n = 3508$) was used for follow-up analyses, where those who were excluded from the LCA ($n = 2663$) are hereafter labelled the Comparison Group. This was the group that did not report a history of suicidal ideation or attempts. Follow-up analyses were conducted in Jamovi (The jamovi project, 2022). Classes were compared at baseline using ANOVAs. Mixed effects logistic regression analysis was used to predict suicide attempts longitudinally. Multinomial regression analyses were used to study class differences at follow-up. More information on each test is available in the supplementary materials.

3. Results

3.1. Description of the latent classes

An overview of the different LCA solutions is reported in Table 1. In the LCA, a three-class solution was selected for the best fit for the current sample ($n = 845$). While the six-class solution had the lowest AIC and BIC values, it had low entropy and various classes smaller than 5 % of the sample. A scree plot which helped to identify which solution to select can be found in Appendix A.

Descriptive statistics by class membership are shown in Table 2. The three classes reflected a group of people with relatively low symptoms (Class 1; $n = 529$, 62 % of the sample), a group of people with moderate symptoms (Class 2; $n = 195$, 23 % of the sample), and a group with high levels of symptoms (Class 3; $n = 119$, 14 % of the sample). These groups were compared against each other, and with the comparison group ($n = 2663$). The groups differed on all variables of interest except age, sex and ethnicity. Full details on the differences between the classes at baseline are in the supplementary materials.

3.2. Predicting suicide attempts at follow up

Out of the 3508 participants, 2418 (68.9 %) completed the 12 months follow up, 2236 (63.7 %) completed the 24 months follow up, and 2226 (63.5 %) completed the third follow-up at 36 months. At 36 months, an ordinal logistic regression revealed significant differences in attrition between the classes. Using the comparison group as a reference group, those in Class 1 were more likely to have missed one or more follow-up, $Wald z = 2.42, p = .015, OR = 1.24$, as well as those in Class 3, $Wald z = 3.42, p < .001, OR = 1.79$. Looking at the demographics, those who remained in the study were slightly older ($M = 25.7, SD = 4.75$) than those who didn't ($M = 25.1, SD = 4.91$), $t(3450) = 3.61, p < .001$, and those who remained were more likely to be female,

$$\chi^2(1) = 84.5, p < .001.$$

Because of this differential dropout, our original analysis plan changed, and the multinomial regression to distinguish between class differences at 36 months is not reported in this paper. However, for transparency, it is reported in the supplementary materials, as we think the results show interesting findings despite the fact that the remaining sample is not representative. Moreover, because of this attrition, a mixed effects logistic regression was used to predict suicide attempts at the three follow-ups. Class assignment was added as a factor, together with Follow-up. This Follow-up variable compared the three different time points at 12-, 24- and 36-months post-baseline. There was a significant effect of Class on the likelihood of a suicide attempt, $F(2, 2451) = 56.27, p < .001$. Post-hoc comparisons showed that there were significant differences between all groups, with the comparison group being the least likely to have a suicide attempt at any of the follow-ups, followed by Class 1, then Class 2, and finally Class 3. We also found a significant effect of Follow-up, $F(2, 4433) = 10.86, p < .001$. There was no significant difference in the likelihood of a suicide attempts across all classes between the 12-month and the 24-month follow-up. There were however significant differences between the 12-month and the 36-month follow-up, $t(4552) = 4.04, p < .001$, indicating that people were more likely to have attempted suicide at 12 months than at 36 months. We found a similar effect for the comparison between 24 months and 36 months, $t(4330) = 4.10, p < .001$. We also found a significant interaction between Follow-up and Class, $F(6, 4390) = 6.65, p < .001$, which we have plotted for ease of interpretation in Fig. 1. We found that the comparison group and Class 1 stayed relatively stable over time, with Class 1 showing a slight decrease by 36 months post-baseline, whereas Class 2 showed a slight increase in the likelihood of a suicide attempt at 24 months compared to 12 months. Class 3 showed the most notable difference, with a much higher likelihood of a suicide attempt at the 12-month follow-up, after which the likelihood decreases with time, albeit still being higher than the other groups.

4. Discussion

The aim of this study was to use Latent Class Analysis (LCA) to identify subgroups within a nationally representative dataset based on established suicide risk factors from the IMV model (O'Connor and Kirtley, 2018). Additionally, we examined how these groups differed at baseline and longitudinally, as well as predicting who was most at risk for future suicidal behaviour. Within a subsample of individuals with a history of suicidal thoughts and/or attempts, we identified three subgroups (based on levels of depression, defeat, internal and external entrapment, suicidal ideation and impulsivity); a low symptom group (Class 1 = 62 %), a moderate symptom group (Class 2 = 23 %), and a high symptom group (Class 3 = 14 %).

Findings suggested that higher levels of symptoms were also associated with higher levels on associated risk variables such as impulsivity and perfectionism, and lower levels of protective variables such as social support. At 36 months follow-up, those in the high symptom class were most likely to have made a suicide attempt in the last 36 months, and the control group having the lowest risk. Comparing the different follow-ups, it seems that the risk of a suicide attempt was highest within 12 months after baseline in the high risk group. However, the moderate risk group showed an increase in suicide attempt rates at the 24-month follow-up, which was retained at 36 months. The low risk group showed a relatively stable risk over time. While a history of suicidal behaviour has been highlighted in many previous studies, both for suicide deaths (Bostwick et al., 2016) and future suicide attempts (O'Connor et al., 2013), entrapment has not had as much attention. Our findings suggest that within a subsample of people with a suicidal history, entrapment may be especially important in distinguishing between those at lower and higher risk for future suicidal behaviour. This finding fits with a clinical prospective study which identified entrapment as a

Table 1
Overview of the solutions from the Latent Class Analysis.

Model	AIC	BIC	Entropy	p-value*
One class	34,287	34,415	1.000	–
Two classes	34,273	34,434	0.538	0.010
Three classes	33,577	33,771	0.888	0.010
Four classes	33,587	33,814	0.655	0.356
Five classes	33,599	33,859	0.590	0.475
Six classes	33,290	33,584	0.640	0.010

* Bootstrapped likelihood ratio test.

Note. values in bold indicate the best solution according to that criterion.

Table 2

Baseline demographic, IMV, and other risk and protective variables among the three latent classes and the control group.

Variable	Control group (n = 2607)	Class 1 (n = 529)	Class 2 (n = 195)	Class 3 (n = 119)	Omnibus F or χ^2	p
Demographics						
Age, M (SD)	25.41 (4.87)	25.62 (4.56)	25.43 (4.75)	25.63 (4.88)	0.35	0.790
Sex (female), n (%)	1293 (49.6)	276 (52.2)	111 (56.9)	53 (44.5)	6.25	0.100
Marital status (married/civil partnership), n (%)	461 (17.7)	79 (14.9)	18 (9.2)	7 (5.9)	20.9	<0.001
Employment status (active) n (%)	2147 (82.8)	382 (72.3)	119 (61.7)	68 (57.6)	105	<0.001
Ethnicity (white) n (%)	2442 (93.7)	500 (94.5)	186 (95.4)	113 (95.0)	1.41	0.703
Sexual orientation (heterosexual) n (%)	1713 (94.6)	319 (90.9)	121 (82.9)	54 (74.0)	69.8	<0.001
Variables for the formation of the latent classes						
BDI, M (SD)	7.59 (7.94)	13.70 (9.46)	29.07 (11.04)	37.16 (11.11)	532.9	<0.001
BSSI 19, M (SD)	0.18 (1.19)	0.67 (1.87)	2.95 (3.59)	16.87 (5.36)	427.6	<0.001
Defeat, M (SD)	12.35 (9.96)	20.62 (11.24)	37.68 (10.87)	45.65 (10.48)	718.4	<0.001
Internal Entrapment, M (SD)	2.16 (3.96)	4.03 (3.66)	17.18 (3.63)	18.93 (4.55)	1463.3	<0.001
External Entrapment, M (SD)	4.08 (6.01)	7.50 (7.06)	20.22 (7.87)	25.82 (7.81)	556.1	<0.001
Impulsivity, M (SD)	60.31 (10.20)	64.89 (10.67)	70.13 (11.33)	72.33 (11.77)	99.8	<0.001
Mental health risk variables						
Social support, M (SD)	26.5 (3.99)	25.0 (4.89)	22.5 (5.06)	19.9 (6.38)	85.7	<0.001
Perceived Burdenomeness, M (SD)	11.7 (5.21)	15.7 (7.08)	25.2 (8.30)	31.7 (8.61)	400.5	<0.001
Thwarted Belongingness, M (SD)	10.2 (5.66)	12.9 (6.48)	18.0 (6.25)	22.6 (6.08)	254.9	<0.001
Socially Prescribed Perfectionism, M (SD)	40.1 (12.49)	44.7 (13.31)	53.5 (12.82)	57.5 (14.20)	125.8	<0.001
ACSS, M (SD)	13.6 (4.10)	14.8 (4.26)	15.1 (4.80)	16.4 (5.11)	25.4	<0.001
Wellbeing, M (SD)	51.9 (7.51)	48.6 (7.35)	39.5 (7.92)	33.9 (7.31)	367.4	<0.001
Problematic alcohol use (yes), n (%)	209 (8.0)	70 (13.2)	36 (18.5)	18 (15.1)	37.2	<0.001
Suicide attempt in the last year (yes), n (%)	0 (0.0)	24 (4.6)	21 (10.8)	43 (36.4)	667	<0.001
History of self-harm (yes), n (%)	165 (6.4)	215 (40.6)	94 (48.2)	76 (63.9)	767	<0.001

Note. Pairwise deletion was used for all analyses, so different total N for each comparison. Full test details can be found in supplementary materials. Bold values indicate significant results. Abbreviations: BDI = Beck Depression Inventory, BSSI = Beck Scale for Suicidal Ideation, ACSS = Acquired Capability for Suicide Scale.




Fig. 1. Plot showing the likelihood of a suicide attempt for each of the follow-ups, separated by Class.

key variable in predicting future suicidal behaviour (O'Connor et al., 2013), as well as Branley-Bell et al. (2019) who called for entrapment to receive closer attention in suicide risk assessments. The current findings are also consistent with the central tenet of the motivational phase of the IMV model, which suggests that entrapment is part of the final common pathway to suicidal thinking and behaviour (O'Connor and Kirtley, 2018). Future research and clinical practice should aim to embed the assessment of entrapment within clinical formulation protocols. As the risk of a suicide attempt was most pronounced in the high risk group at 12 months, and decreased after that, it highlights that long-term prediction may be less viable when assessing suicide risk.

We were surprised that the classes did not differ from each other on acquired capability as the IPTS suggests that those at higher risk of attempting suicide have a higher capability for suicide (Joiner, 2005).

However, our findings are more consistent with Ma et al. (2019) who did not find differences in capability for suicide in their group analysis (Ma et al., 2019). One possible explanation for our null finding is that all of the groups contained some individuals with a previous suicide attempt, thereby rendering it difficult to detect differences in capability between the classes. While this study grouped together those who simply think about suicide and those who attempt suicide, which some argue are two categorically distinct groups (Witte et al., 2017), it does show that high-risk individuals are more likely to attempt suicide at a later point and that these individuals differ on some key components of the ideation-to-action framework (David Klonsky et al., 2017) from those who are at a lower risk, both at baseline and longitudinally.

5. Limitations

It is important to acknowledge our study limitations. First, in the prospective analyses, it is likely that our findings have been affected by differential drop-out as those in the high symptom class were most likely to have dropped out of the study, leading to this group being under-represented in the follow-up analysis. This dropout prevented us from drawing many meaningful conclusions about the stability of class membership over time, which was one of our original aims. However, we do believe that the remaining results are still important additions to the suicide literature. Second, like many similar studies, as all measures were self-report, the findings may be affected by demand characteristics and the under-reporting of sensitive issues such as suicidal thoughts and behaviours. Lastly, despite being a nationally representative sample, the total number of suicide attempts reported by participants was low, which may limit the statistical power. In future research, larger or more high-risk samples should be considered. It is also unclear the extent to which these findings generalise to other countries, in particular those from other cultures. Lastly, we were unable to use longitudinal models such as Latent Class Growth Modelling (LCGM) to investigate changes in symptoms over time. Future research should aim to collect the same variables at baseline and follow-up in order to allow for these more sophisticated methods of predicting suicide-related outcomes. This would allow for studying trajectories of symptom severity related to suicide risk over time and identifying key constructs that affect these trajectories.

6. Conclusions

Overall, our findings emphasise the importance of previous suicidal behaviour and entrapment when investigating suicide risk. To our knowledge, this is the first study using constructs from the IMV model to create latent classes to predict future suicidal behaviour. It provides a solid basis for future research in this area. These findings highlight the potential utility of latent classes in the understanding and prediction of suicide risk.

Declaration of competing interest

None.

Acknowledgements

None.

Financial support

This study was originally funded by a grant from US Department of Defense (W81XWH-12-1-0007). Further funding was provided by a grant from the Mindstep Foundation. The funders had no role in study design, data collection, data analysis, data interpretation or writing of this article.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jad.2023.05.028>.

References

Allan, N.P., Holm-Denoma, J., Conner, K.R., Zuromski, K.L., Saulnier, K.G., Stecker, T., 2020. Profiles of risk for suicidal behavior in past and current United States military personnel: latent profile analysis of current risk factors. *Arch. Suicide Res.* 24 (1), 1–17. <https://doi.org/10.1080/13811118.2018.1506843>.

Au, J.S., Martinez de Andino, A., Mekawi, Y., Silverstein, M.W., Lamis, D.A., 2021. Latent class analysis of bipolar disorder symptoms and suicidal ideation and behaviors. *Bipolar Disord.* 23 (2), 186–195. <https://doi.org/10.1111/bdi.12967>.

Beck, A.T., Steer, R.A., Ranieri, W.F., 1988. Scale for suicide ideation: psychometric properties of a self-report version. *J. Clin. Psychol.* 44 (4), 499–505.

Beck, A.T., Steer, R.A., Brown, G.K., 1996. Beck Depression Inventory Manual. The Psychological Corporation, San Antonio, TX.

Bernanke, J., Galfalvy, H.C., Mortali, M.G., Hoffman, L.A., Moutier, C., Nemeroff, C.B., Stanley, B.H., Clayton, P., Harkavy-Friedman, J., Oquendo, M.A., 2017. Suicidal ideation and behavior in institutions of higher learning: a latent class analysis. *J. Psychiatr. Res.* 95, 253–259. <https://doi.org/10.1016/j.jpsychires.2017.09.003>.

Bostwick, J.M., Pabbati, C., Geske, J.R., McKeon, A.J., 2016. Suicide attempt as a risk factor for completed suicide: even more lethal than we knew. *Am. J. Psychiatry* 173, 1094–1100. <https://doi.org/10.1176/appi.ajp.2016.15070854>.

Branley-Bell, D., O'Connor, D.B., Green, J.A., Ferguson, E., O'Carroll, R.E., O'Connor, R.C., 2019. Distinguishing suicide ideation from suicide attempts: further test of the integrated motivational-volitional model of suicidal behaviour. *J. Psychiatr. Res.* 117, 100–107. <https://doi.org/10.1016/j.jpsychires.2019.07.007>.

David Klonsky, E., Qiu, T., Saffer, B.Y., 2017. Recent advances in differentiating suicide attempts from suicide ideators. *Curr. Opin. Psychiatry* 30 (1), 15–20. <https://doi.org/10.1097/YCO.0000000000000294>.

Dhingra, K., Boduszek, D., O'Connor, R.C., 2015. Differentiating suicide attempts from suicide ideators using the integrated motivational-volitional model of suicidal behaviour. *J. Affect. Disord.* 186, 211–218. <https://doi.org/10.1016/j.jad.2015.07.007>.

Ewing, J.A., 1984. Detecting alcoholism: the CAGE questionnaire. *JAMA J. Am. Med. Assoc.* 252, 1905–1907.

Franklin, J.C., Ribeiro, J.D., Fox, K.R., Bentley, K.H., Kleiman, E.M., Huang, X., Musacchio, K.M., Jaroszewski, A.C., Chang, B.P., Nock, M.K., 2017. Risk factors for suicidal thoughts and behaviors: a meta-analysis of 50 years of research. *Psychol. Bull.* 143 (2), 187–232. <https://doi.org/10.1037/bul0000084>.

Giano, Z., Currin, J.M., Deboy, K., Hubach, R.D., 2020. Identifying distinct subgroups of lesbian, gay, and bisexual youth for suicide risk: a latent profile analysis. *J. Adolesc. Health* 67 (2), 194–200. <https://doi.org/10.1016/j.jadohealth.2020.03.031>.

Gilbert, P., Allan, S., 1998. The role of defeat and entrapment (arrested flight) in depression: an exploration of an evolutionary view. *Psychol. Med.* 28 (3), 585–598.

Ginley, M.K., Bagge, C.L., 2017. Psychiatric heterogeneity of recent suicide attempts: a latent class analysis. *Psychiatry Res.* 251 (November 2016), 1–7. <https://doi.org/10.1016/j.psychires.2017.02.004>.

Hewitt, P.L., Flett, G.L., 1991. Perfectionism in the self and social contexts: conceptualization, assessment, and association with psychopathology. *J. Pers. Soc. Psychol.* 60 (3), 456–470. <https://doi.org/10.1037/0022-3514.60.3.456>.

Hoogstoel, F., Samadoulougou, S., Lorant, V., Kirakoya-Samadoulougou, F., 2021. A latent class analysis of health lifestyles in relation to suicidality among adolescents in Mauritius. *Int. J. Environ. Res. Public Health* 18 (13), 6934. <https://doi.org/10.3390/ijerph18136934>.

IBM Corp., 2015. IBM SPSS Statistics for Windows (No. 23). IBM Corp.

Jiang, Y., Perry, D.K., Hesser, J.E., 2010. Suicide patterns and association with predictors among Rhode Island public high school students: a latent class analysis. *Am. J. Public Health* 100 (9), 1701–1707. <https://doi.org/10.2105/AJPH.2009.183483>.

Joiner, T.E., 2005. Why people die by suicide. In: Why People Die by Suicide. Harvard University Press. <https://doi.org/10.2307/j.ctvjqhv2f>.

Jung, S., Lee, D., Park, S., Hong, H.J., 2019. Subtypes of suicidal ideation in Korean adolescents: a multilevel latent profile analysis. *Aust. N. Z. J. Psychiatry* 53 (2), 158–167. <https://doi.org/10.1177/0004867418763530>.

Logan, J., Hall, J., Karch, D., 2011. Suicide categories by patterns of known risk factors: a latent class analysis. *Arch. Gen. Psychiatry* 68 (9), 935–941. <https://doi.org/10.1001/archgenpsychiatry.2011.85>.

Love, H.A., Durtschi, J.A., 2021. Suicidal ideation and behaviors in young adults: a latent profile analysis. *J. Fam. Psychol.* 35 (3), 345–355. <https://doi.org/10.1037/fam0000786>.

Lucht, L., Höller, I., Forkmann, T., Teismann, T., Schönfelder, A., Rath, D., Paashaus, L., Stengler, K., Juckel, G., Glaesmer, H., 2020. Validation of the motivational phase of the integrated motivational-volitional model of suicidal behavior in a German high-risk sample. *J. Affect. Disord.* 274, 871–879. <https://doi.org/10.1016/j.jad.2020.05.079>.

Ma, J.S., Batterham, P.J., Calear, A.L., Han, J., 2019. Suicide risk across latent class subgroups: a test of the generalizability of the interpersonal psychological theory of suicide. *Suicide Life Threat. Behav.* 49 (1), 137–154. <https://doi.org/10.1111/sltb.12426>.

McManus, S., Meltzer, H., Brugha, T., Bebbington, P.E., Jenkins, R., 2009. Adult Psychiatric Morbidity in England: Results of a Household Survey. <https://doi.org/10.13140/2.1.1563.5205>.

Mitchell, P.H., Powell, L., Blumenthal, J., Norten, J., Ironson, G., Pitula, C.R., Froelicher, E.S., Czajkowski, S., Youngblood, M., Huber, M., others, 2003. A short social support measure for patients recovering from myocardial infarction: the ENRICHD social support inventory. *J. Cardiopulm. Rehab. Prev.* 23 (6), 398–403.

O'Connor, R.C., Kirtley, O.J., 2018. The integrated motivational-volitional model of suicidal behaviour. *Philos. Trans. R. Soc. Ser. B Biol. Sci.* 373 (1754) <https://doi.org/10.1098/rstb.2017.0268>.

O'Connor, R.C., Smyth, R., Ferguson, E., Ryan, C., Williams, J.M.G., 2013. Psychological processes and repeat suicidal behavior: a four-year prospective study. *J. Consult. Clin. Psychol.* 81 (6), 1137–1143. <https://doi.org/10.1037/a0033751>.

O'Connor, R.C., Wetherall, K., Cleare, S., Eschle, S., Drummond, J., Ferguson, E., O'Connor, D.B., O'Carroll, R.E., 2018. Suicide attempts and non-suicidal self-harm: national prevalence study of young adults. *BJPsych Open* 4 (3), 142–148. <https://doi.org/10.1192/bjop.2018.14>.

O'Connor, D.B., Branley-Bell, D., Green, J., Ferguson, E., O'Carroll, R., O'Connor, R.C., 2021. Resilience and vulnerability factors influence daily cortisol levels in individuals vulnerable to suicide. *J. Psychiatr. Res.* 142, 312–320.

O'Neill, S., McLafferty, M., Ennis, E., Lapsley, C., Bjourson, T., Armour, C., Murphy, S., Bunting, B., Murray, E., 2018. Socio-demographic, mental health and childhood adversity risk factors for self-harm and suicidal behaviour in College students in Northern Ireland. *J. Affect. Disord.* 239, 58–65. <https://doi.org/10.1016/j.jad.2018.06.006>.

Ordóñez-Carrasco, J.L., Cuadrado-Guirado, I., Rojas-Tejada, A.J., 2020. Frustrated interpersonal needs as a motivational moderator in the integrated motivational-volitional model. *Death Stud.* <https://doi.org/10.1080/07481187.2020.1783031>.

Patton, J.H., Stanford, M.S., Barratt, E.S., 1995. Factor structure of the Barratt impulsiveness scale. *J. Clin. Psychol.* 51 (6), 768–774.

Rice, S.M., Oliiffe, J.L., Kealy, D., Ogorodniczuk, J.S., 2018. Male depression subtypes and suicidality latent profile analysis of internalizing and externalizing symptoms in a representative Canadian sample. *J. Nerv. Ment. Dis.* 206 (3), 169–172. <https://doi.org/10.1097/NMD.0000000000000739>.

Rosenberg, J.M., Beymer, P.N., Anderson, D.J., Van Lissa, C.J., Schmidt, J.A., 2018. tidyLPA: an R Package to easily carry out latent profile analysis (LPA) using open-source or commercial software. *J. Open Source Softw.* 3 (30), 978. <https://doi.org/10.21105/joss.00978>.

Selby, E.A., Gardner, K., 2015. A latent profile analysis of suicidal and self-injurious behavior, other dysregulated behaviors, and borderline personality disorder symptoms. *J. Exp. Psychopathol.* 6 (4), 356–368. <https://doi.org/10.5127/jep.034913>.

Tennant, R., Hiller, L., Fishwick, R., Platt, S., Joseph, S., Weich, S., Parkinson, J., Secker, J., Stewart-Brown, S., 2007. The Warwick-Edinburgh mental well-being scale (WEMWBS): development and UK validation. *Health Qual. Life Outcomes* 5 (1), 63. <https://doi.org/10.1186/1477-7525-5-63>.

The jamovi project, 2022. *jamovi* (Version 2.3) [Computer Software]. Retrieved from. <https://www.jamovi.org>.

Thullen, M.J., Taliaferro, L.A., Muehlenkamp, J.J., 2016. Suicide ideation and attempts among adolescents engaged in risk behaviors: a latent class analysis. *J. Res. Adolesc.* 26 (3), 587–594. <https://doi.org/10.1111/jora.12199>.

van Orden, K.A., Witte, T.K., Gordon, K.H., Bender, T.W., Joiner, T.E., 2008. Suicidal desire and the capability for suicide: tests of the interpersonal-psychological theory of suicidal behavior among adults. *J. Consult. Clin. Psychol.* 76 (1), 72–83. <https://doi.org/10.1037/0022-006X.76.1.72>.

van Orden, K.A., Cukrowicz, K.C., Witte, T.K., Joiner, T.E., 2012. Thwarted belongingness and perceived burdensomeness: construct validity and psychometric properties of the Interpersonal Needs Questionnaire. *Psychol. Assess.* 24 (1), 197–215. <https://doi.org/10.1037/a0025358>.

Wetherall, Karen, Cleare, S., Eschle, S., Ferguson, E., O'Connor, D.B., O'Carroll, R.E., O'Connor, R.C., 2018. From ideation to action: differentiating between those who think about suicide and those who attempt suicide in a national study of young adults. *J. Affect. Disord.* 241 (June), 475–483. <https://doi.org/10.1016/j.jad.2018.07.074>.

Wetherall, K., Cleare, S., Zortea, T., O'Connor, R., 2020. Status of the integrated motivational-volitional model of suicidal behavior. In: de Leo, D., Postuvan, V. (Eds.), *Reducing the Toll of Suicide: Resources for Communities, Groups, and Individuals*. Hogrefe Publishing, pp. 169–184. <https://psycnet.apa.org/record/2019-67864-011>.

Witte, T.K., Holm-Denoma, J.M., Zuromski, K.L., Gauthier, J.M., Ruscio, J., 2017. Individuals at high risk for suicide are categorically distinct from those at low risk. *Psychol. Assess.* 29 (4), 382–393. <https://doi.org/10.1037/pas0000349>.

Wong, Q.J.J., Torok, M., van Spijk, B.A.J., Werner-Seidler, A., Calear, A.L., Batterham, P.J., Han, J., Christensen, H., 2020. Identifying subgroups within a sample of adults with a suicide attempt history using the Interpersonal Psychological Theory of Suicide. *Psychiatry Res.* 293 (May), 113406. <https://doi.org/10.1016/j.psychres.2020.113406>.

World Health Organization (WHO), 2021. Suicide worldwide in 2019: global health estimates. <https://apps.who.int/iris/bitstream/handle/10665/341728/9789240026643-eng.pdf?sequence=1>.

Xiao, Y., Romanelli, M., Lindsey, M.A., 2019. A latent class analysis of health lifestyles and suicidal behaviors among US adolescents. *J. Affect. Disord.* 255 (May), 116–126. <https://doi.org/10.1016/j.jad.2019.05.031>.

Zhu, C., Chen, L., Ou, L., Geng, Q., Jiang, W., Lv, X., Wu, X., Ci, H., Liu, Q., Yao, Y., Pentadibiran, P., Persekutuan, K., Kami, R., Ketua, S., Kementerian, S., Persekutuan, J., Pentadibiran, S., Kerajaan, S., Berkanun, B., Flynn, D., 2019. Suicidal ideation in emerging and young adults: latent profile analysis of risk and protective factors in a nationally representative sample, 8 (2). <https://doi.org/10.22201/fq.18708404e.2004.3.66178>, 2019.